

Mouse Free Fatty Acid Receptor 2 (FFAR2) ELISA Kit | FFAR2 elisa kit

Mouse Free Fatty Acid Receptor 2 (FFAR2) ELISA Kit

CATALOG #
AAA67293

REACTIVITY
Mouse


PRODUCT SYNONYMS

Free Fatty Acid Receptor 2 (FFAR2); Mouse Free Fatty Acid Receptor 2 (FFAR2) ELISA Kit; FFAR2 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of FFAR2. No significant cross-reactivity or interference between FFAR2 and analogues was observed.
Sequence Length	330
Samples	Mouse tissue homogenates, cell lysates or other biological fluids
Assay Type	Quantitative Sandwich
Detection Range	0.156-10ng/mL
Sensitivity	< 0.055ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level FFAR2 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level FFAR2 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for FFAR2 elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for the in vitro quantitative measurement of FFAR2 in mouse tissue homogenates, cell lysates or other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to FFAR2. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to FFAR2. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain FFAR2, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of FFAR2 in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	270341373
NCBI GeneID	233079
NCBI Accession #	NP_001161981.1
NCBI GenBank Nucleotide #	NM_001168509.1
UniProt Accession #	Q8VCK6
Molecular Weight	37,124 Da
NCBI Official Full Name	free fatty acid receptor 2
NCBI Official Synonym Full Names	free fatty acid receptor 2
NCBI Official Symbol	Ffar2
NCBI Official Synonym Symbols	Gpr43; GPCR43
NCBI Protein Information	free fatty acid receptor 2
UniProt Protein Name	Free fatty acid receptor 2
UniProt Gene Name	Ffar2
UniProt Synonym Gene Names	Gpr43; Lssig

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/mouse-elisa-kits/free-fatty-acid-receptor-2-ffar2/67293>