

Human Myostatin (MSTN) ELISA Kit | MSTN elisa kit

Human Myostatin (MSTN) ELISA Kit

CATALOG #
AAA67441

REACTIVITY
Human


PRODUCT SYNONYMS

Myostatin (MSTN); Human Myostatin (MSTN) ELISA Kit; GDF8 Growth Differentiation Factor 8; MSTN elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of MSTN. No significant cross-reactivity or interference between MSTN and analogues was observed.
Sequence Length	375
Samples	Serum, plasma, tissue homogenates, cell lysates, cell culture supernates or other biological fluids
Assay Type	Quantitative Sandwich
Detection Range	0.781-50ng/mL
Sensitivity	< 0.25ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level MSTN were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level MSTN were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100.
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for MSTN elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for the in vitro quantitative measurement of MSTN in human serum, plasma, tissue homogenates, cell lysates, cell culture supernates or other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to MSTN. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to MSTN. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain MSTN, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of MSTN in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	4885259
NCBI GeneID	2660
NCBI Accession #	NP_005250.1
NCBI GenBank Nucleotide #	NM_005259.2
UniProt Accession #	O14793; Q6B0H2; A1C2J7; A1C2K0
Molecular Weight	42,750 Da
NCBI Official Full Name	growth/differentiation factor 8 preproprotein
NCBI Official Synonym Full Names	myostatin
NCBI Official Symbol	MSTN
NCBI Official Synonym Symbols	GDF8; MSLHP
NCBI Protein Information	growth/differentiation factor 8
UniProt Protein Name	Growth/differentiation factor 8
UniProt Gene Name	MSTN
UniProt Synonym Gene Names	GDF8; GDF-8

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.