

Canine Gastrin (GT) ELISA Kit | GT elisa kit

Canine Gastrin (GT) ELISA Kit

CATALOG #
AAA67700

REACTIVITY
Canine


PRODUCT SYNONYMS

Gastrin (GT); Canine Gastrin (GT) ELISA Kit; GT elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Canine
Specificity	This assay has high sensitivity and excellent specificity for detection of GT. No significant cross-reactivity or interference between GT and analogues was observed.
Sequence Length	104
Samples	Serum, plasma and other biological fluids.
Assay Type	Competitive inhibition
Detection Range	15.625-1000pg/mL
Sensitivity	4.28pg/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level GT were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level GT were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for GT elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of GT in canine serum, plasma and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to GT. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to GT. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain GT, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of GT in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	345805367
NCBI GeneID	100685087
NCBI Accession #	XP_003435286.1
NCBI GenBank Nucleotide #	XP_003435286.1
UniProt Accession #	P01353
Molecular Weight	11,519 Da
NCBI Official Full Name	gastrin isoform X2
NCBI Official Symbol	GAST
NCBI Official Synonym Symbols	GAS
NCBI Protein Information	gastrin
UniProt Protein Name	Gastrin
UniProt Gene Name	GAST
UniProt Synonym Gene Names	GAS; G34

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.