

Human Interleukin 16 (IL16) ELISA Kit | IL16 elisa kit

Human Interleukin 16 (IL16) ELISA Kit

CATALOG #
AAA67743

REACTIVITY
Human


PRODUCT SYNONYMS

Interleukin 16 (IL16); Human Interleukin 16 (IL16) ELISA Kit; LCF; PrIL-16; Lymphocyte Chemoattractant Factor; Pro-Interleukin 16; IL16 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of IL16. No significant cross-reactivity or interference between IL16 and analogues was observed.
Sequence Length	1331
Samples	Serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids.
Assay Type	Sandwich
Detection Range	15.625-1000pg/mL
Sensitivity	6.1pg/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level IL16 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level IL16 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for IL16 elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of IL16 in human serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to IL16. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to IL16. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain IL16, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of IL16 in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	289063395
NCBI GeneID	3603
NCBI Accession #	NP_001165599.1
NCBI GenBank Nucleotide #	NM_001172128.1
UniProt Accession #	Q14005; Q16435; Q6VVE6; Q6ZMQ7; Q9UP18; A6NM20; A8MU65; B5TY35; B9EGR6; H3BVH5
Molecular Weight	59,345 Da
NCBI Official Full Name	pro-interleukin-16 isoform 3
NCBI Official Synonym Full Names	interleukin 16
NCBI Official Symbol	IL16
NCBI Official Synonym Symbols	LCF; NIL16; PRIL16; prIL-16
NCBI Protein Information	pro-interleukin-16
UniProt Protein Name	Pro-interleukin-16
UniProt Gene Name	IL16
UniProt Synonym Gene Names	IL-16; LCF

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.