

Human Transferrin Receptor (TFR) ELISA Kit | TFR elisa kit

Human Transferrin Receptor (TFR) ELISA Kit

CATALOG #
AAA67861

REACTIVITY
Human


PRODUCT SYNONYMS

Transferrin Receptor (TFR); Human Transferrin Receptor (TFR) ELISA Kit; TFR elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of TFR. No significant cross-reactivity or interference between TFR and analogues was observed.
Sequence Length	760
Samples	Serum, plasma, tissue homogenates, cell lysates and other biological fluids.
Assay Type	Sandwich
Detection Range	1.56-100ng/mL
Sensitivity	0.58ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level TFR were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level TFR were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for TFR elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of TFR in human serum, plasma, tissue homogenates, cell lysates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to TFR. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to TFR. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain TFR, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of TFR in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	189458819
NCBI GeneID	7037
NCBI Accession #	NP_001121620.1
NCBI GenBank Nucleotide #	NM_001128148.2
UniProt Accession #	P02786; Q1HE24; Q59G55; Q9UCN0; Q9UCU5; Q9UDF9; Q9UK21; D3DXB0
Molecular Weight	84,871 Da
NCBI Official Full Name	transferrin receptor protein 1 isoform 1
NCBI Official Synonym Full Names	transferrin receptor
NCBI Official Symbol	TFRC
NCBI Official Synonym Symbols	T9; TR; TFR; p90; CD71; TFR1; TRFR; IMD46
NCBI Protein Information	transferrin receptor protein 1
UniProt Protein Name	Transferrin receptor protein 1
UniProt Gene Name	TFRC
UniProt Synonym Gene Names	TR; TfR; TfR1; Trfr; sTfR

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.