

Mouse Aromatase (ARO) ELISA Kit | ARO elisa kit

Mouse Aromatase (ARO) ELISA Kit

CATALOG # AAA68501	REACTIVITY Mouse
------------------------------	----------------------------


PRODUCT SYNONYMS

Aromatase (ARO); Mouse Aromatase (ARO) ELISA Kit; CYP19A1; CYPXIX; ARO; ARO1; CPV1; CYAR; CYP19; P450AROM; P450-AROM; Estrogen synthase; Cytochrome P450; Family 19; Subfamily A; Polypeptide 1; ARO elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of ARO. No significant cross-reactivity or interference between ARO and analogues was observed.
Sequence Length	503
Samples	Serum, plasma, tissue homogenates or other biological fluids
Assay Type	Quantitative Sandwich
Detection Range	0.312-20ng/mL
Sensitivity	<0.117ng/mL.
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level ARO were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level ARO were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for ARO elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for the in vitro quantitative measurement of ARO in mouse serum, plasma, tissue homogenates or other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to ARO. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to ARO. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain ARO, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of ARO in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	1131345323
NCBI GeneID	13075
NCBI Accession #	NP_001335100.1
NCBI GenBank Nucleotide #	NM_001348171.1
UniProt Accession #	P28649
Molecular Weight	58,015 Da
NCBI Official Full Name	aromatase
NCBI Official Synonym Full Names	cytochrome P450, family 19, subfamily a, polypeptide 1
NCBI Official Symbol	Cyp19a1
NCBI Official Synonym Symbols	Ar; ArKO; Int5; Cyp19; Int-5; p450arom
NCBI Protein Information	aromatase
UniProt Protein Name	Aromatase
UniProt Gene Name	Cyp19a1
UniProt Synonym Gene Names	Arom; Cyp19

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.