

Human Transient Receptor Potential Cation Channel Subfamily M, Member 7 (TRPM7) ELISA Kit | TRPM7 elisa kit

Human Transient Receptor Potential Cation Channel Subfamily M, Member 7 (TRPM7) ELISA Kit

CATALOG # AAA68737	REACTIVITY Human
------------------------------	----------------------------


PRODUCT SYNONYMS

Transient Receptor Potential Cation Channel Subfamily M; Member 7 (TRPM7); Human Transient Receptor Potential Cation Channel Subfamily M; Member 7 (TRPM7) ELISA Kit; CHAK; CHAK1; LTRPC7; TRP-PLIK; Channel-kinase 1; Long transient receptor potential channel 7; TRPM7 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of TRPM7. No significant cross-reactivity or interference between TRPM7 and analogues was observed.
Sequence Length	1864
Samples	Human tissue homogenates or other biological fluids
Assay Type	Quantitative Sandwich
Detection Range	78.1-5000pg/mL
Sensitivity	< 33pg/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level TRPM7 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level TRPM7 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for TRPM7 elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for the in vitro quantitative measurement of TRPM7 in human tissue homogenates or other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to TRPM7. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to TRPM7. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain TRPM7, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of TRPM7 in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	667751393
NCBI GeneID	54822
NCBI Accession #	NP_001288141.1
NCBI GenBank Nucleotide #	NM_001301212.1
UniProt Accession #	Q96QT4; Q6ZMF5; Q86VJ4; Q8NBW2; Q9BXB2; Q9NXQ2
Molecular Weight	212,697 Da
NCBI Official Full Name	transient receptor potential cation channel subfamily M member 7 isoform b
NCBI Official Synonym Full Names	transient receptor potential cation channel subfamily M member 7
NCBI Official Symbol	TRPM7
NCBI Official Synonym Symbols	CHAK; CHAK1; ALSPDC; LTRPC7; LTrpC-7; TRP-PLIK
NCBI Protein Information	transient receptor potential cation channel subfamily M member 7
UniProt Protein Name	Transient receptor potential cation channel subfamily M member 7
UniProt Gene Name	TRPM7
UniProt Synonym Gene Names	CHAK1; LTRPC7; LTrpC-7; LTrpC7

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/human-elisa-kits/transient-receptor-potential-cation-channel-subfamily-m-member-7-trpm7/68737>