

anti-Human TIGIT Antibody | anti-TIGIT antibody

Anti-TIGIT [4A5]

CATALOG # AAA72361	REACTIVITY Human	APPLICATIONS Flow Cytometry, Functional Assay, ELISA
------------------------------	----------------------------	--


PRODUCT SYNONYMS

TIGIT; Anti-TIGIT [4A5]; VSTM3; VSIG9; WUCAM; T-cell immunoreceptor with Ig and ITIM domains; V-set and immunoglobulin domain-containing protein 9; V-set and transmembrane domain-containing protein 3; anti-TIGIT antibody


PRODUCT SPECIFICATIONS

Reactivity	Human
Isotype	Rabbit IgG kappa
Clone Number	4A5
Specificity	This antibody binds the extracellular domain of the human TIGIT (T cell immunoreceptor with Ig and ITIM domains). TIGIT is a type I transmembrane protein that is a member of the CD28 family within the Ig superfamily of proteins. It is an immune receptor present on some T cells and natural killer cells (NK). It binds the poliovirus receptor (PVR) with high affinity which causes increased secretion of IL10 and decreased secretion of IL12B and suppresses T-cell activation by promoting the generation of mature immunoregulatory dendritic cells.
Purity/Purification	Purified. Protein A affinity purified
Form/Format	PBS with 0.02% Proclin 300.
Concentration	1mg/ml (varies by lot)
Applicable Applications for anti-TIGIT antibody	Blocking, FCM/FACS (Flow Cytometry), ELISA
Preparation and Storage	Store at 4 degree C for up to 3 months. For longer storage, aliquot and store at -20 degree C.

IMAGES



IF (Immunofluorescence)
Immunofluorescence staining of human peripheral blood leucocytes with anti-TIGIT antibody 4A5 . Immunofluorescence analysis of paraformaldehyde fixed human peripheral blood leucocytes on Shi-fix™ coverslips stained with the chimeric rabbit IgG version of 4A5 (AAA72361) (1:100 dilution) for 1h followed by Alexa Fluor 488 secondary antibody (1:1000 dilution), showing membrane and cytoplasmic staining. The nuclear stain is DAPI (blue). Panels show, from left-right, top-bottom, AAA72361, DAPI, merged channels, and an isotype control. The isotype control was an unknown specificity antibody followed by staining with Alexa Fluor 488 secondary antibody.



WB (Western Blot)
Western blot using anti-TIGIT antibody 4A5 . Human spleen (A), tonsil (B), and lymph node (C) tissue lysates (35ug protein in RIPA buffer) were resolved on an SDS-PAGE gel, and blots were probed with the chimeric rabbit version of 4A5 (AAA72361) at 1ug/ml before detection using an anti-rabbit secondary antibody. A primary incubation of 1h was used, and protein was detected by chemiluminescence.

ADDITIONAL INFORMATION

Related Product Information for anti-TIGIT antibody	The original antibody was generated by immunizing 9 week old BALB/c mice with recombinant human TIGIT extracellular region Fc fusion protein in complete Freund's adjuvant. This chimeric rabbit antibody was made using the variable domain sequences of the original Mouse IgG1 format for improved compatibility with existing reagents assays and techniques.
Product Categories/Family for anti-TIGIT antibody	cancer; oncology; HIV infection; T-cell mediated immunity

NCBI AND UNIPROT INFORMATION

NCBI GI #	121943253
NCBI GeneID	201633
UniProt Accession #	Q495A1; Q495A3; Q5JPD8; Q6MZS2; Q8N877
Molecular Weight	18,432 Da
NCBI Official Full Name	T-cell immunoreceptor with Ig and ITIM domains
NCBI Official Synonym Full Names	T cell immunoreceptor with Ig and ITIM domains
NCBI Official Symbol	TIGIT
NCBI Official Synonym Symbols	VSIG9; VSTM3; WUCAM
NCBI Protein Information	T-cell immunoreceptor with Ig and ITIM domains
UniProt Protein Name	T-cell immunoreceptor with Ig and ITIM domains
UniProt Gene Name	TIGIT
UniProt Synonym Gene Names	VSIG9; VSTM3

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

