

## Human Dengue Virus IgG ELISA Kit

Human Dengue Virus IgG ELISA Kit

CATALOG #  
**AAA72643**

REACTIVITY  
**Human**

### PRODUCT SYNONYMS

Dengue Virus IgG; Human Dengue Virus IgG ELISA Kit; Dengue Virus IgG elisa kit

### PRODUCT SPECIFICATIONS

|                         |                                                                                |
|-------------------------|--------------------------------------------------------------------------------|
| Reactivity              | Human                                                                          |
| Samples                 | Human Serum, Plasma, Body Fluids, Tissue Homogenate Or Cell Culture Supernates |
| Assay Type              | Sandwich                                                                       |
| Detection Range         | 312 pg/ml-20,000 pg/ml                                                         |
| Sensitivity             | < 10 pg/ml                                                                     |
| Preparation and Storage | Storage 2 - 8 degree C                                                         |

### ADDITIONAL INFORMATION

#### Related Product Information for Dengue Virus IgG elisa kit

Intended Uses: The Cystatin C ELISA has been designed for the precise measurement of Cystatin C in human serum, plasma, body fluids, tissue homogenate or cell culture supernates. For research use only.

Principle of the Assay: This kit is based on a sandwich enzyme-linked immune-sorbent assay technology. An analyte-specific polyclonal antibody is pre-coated onto 96-well plates. The biotin conjugated second antibody is used as detection antibody. During the first incubation the standards and samples react with the analyte-specific pre-coated antibody. In a second incubation the biotin conjugated detection antibody completes the sandwich. After washing with wash buffer the TMB substrate is added to visualize HRP enzymatic reaction. TMB is catalyzed by HRP to produce a blue color product that changes into yellow after adding acidic stop solution. The optical density of the yellow color is proportional to the analyte captured in plate. Read the O.D. absorbance at 450nm in a microtiterplate reader, and calculate the concentration of the analyte in the sample by taking into consideration the dilution factor of the sample!!Background/Introduction: Cystatin C or cystatin 3, a protein encoded by the CST3 gene, is mainly used as a biomarker of kidney function. It is a single 120-residue polypeptide with a molecular mass of approximately 13.26 kD. Cystatin C plays a role in network reorganization in the epileptic dentate gyrus, especially in granule cell dispersion and guidance of migrating newborn granule cells. It is an alternative measure of kidney function that may have prognostic importance among elderly persons who do not meet standard criteria for chronic kidney disease. Among elderly persons without chronic kidney disease, Cystatin C is a prognostic biomarker of risk for death, cardiovascular disease, and chronic kidney disease, thereby signaling a state of 'preclinical' kidney dysfunction.

**For Research Use Only.** Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | [contact@aaabiotech.com](mailto:contact@aaabiotech.com)

Online datasheet: <https://www.aaabiotech.com/human-elisa-kits/dengue-virus-igg/72643>