

MARK2 recombinant protein

MARK2, Active

CATALOG # AAA73679	HOST Sf9 insect cells	APPLICATIONS Western Blot
------------------------------	---------------------------------	-------------------------------------


PRODUCT SYNONYMS

MARK2; Active; EMK1; PAR-1; MGC99619; MARK2 recombinant protein

PRODUCT SPECIFICATIONS


Host	Sf9 insect cells
Form/Format	Recombinant protein stored in 50mM Tris-HCl, pH7.5, 150mM NaCl, 10mM glutathione, 0.1mM EDTA, 0.25mM DTT, 0.1mM PMSF, 25% glycerol.
Applicable Applications for MARK2 recombinant protein	WB (Western Blot), Kinase Assay
Tag	GST
Species	Human
Dry Ice Note	The product may be shipped with dry ice, and an additional fee may be added to your shipping cost.
Preparation and Storage	Store product at -70 degree C. For optimal storage, aliquot target into smaller quantities after centrifugation and store at recommended temperature. For most favorable performance, avoid repeated handling and multiple freeze/thaw cycles.

IMAGES



SDS-PAGE

Purity: Sample Purity Data. For specific information on a given lot, see related technical data sheet.



Activity

Specific Activity: Sample Kinase Activity Plot. For specific information on a given lot, see related technical data sheet.

ADDITIONAL INFORMATION

Related Product Information for MARK2 recombinant protein	Recombinant full-length human MARK2 was expressed by baculovirus in Sf9 insect cells using an N-terminal GST tag. The MARK2 gene accession number is NM_001039469. MARK2 or microtubule affinity-regulating kinase 2 is a member of the Par-1 family of serine/threonine protein kinases. MARK2 is an important regulator of cell polarity in epithelial and neuronal cells and controls the stability of microtubules through phosphorylation and inactivation of several microtubule-associating proteins (1). MARK2 is a key target of H. pylori CagA in the disorganization of gastric epithelial architecture underlying mucosal damage, inflammation, and carcinogenesis (2). MARK2 phosphorylates kinesin-like motor protein GAKIN/KIF13B to regulate axon formation. MARK2 function in the establishment of T cell polarity following engagement to an APC.
Product Categories/Family for MARK2 recombinant protein	Protein Kinases; Invasion/Metastasis; Neurobiology; Ser/Thr Kinases

References

1. Yoshimura, Y. et.al: Polarity-regulating kinase partitioning-defective 1/microtubule affinity-regulating kinase 2 negatively regulates development of dendrites on hippocampal neurons.. J. Neurosci. 27:13098-13107, 2010
2. Saadat, I. et.al: Helicobacter pylori CagA targets PAR1/MARK kinase to disrupt epithelial cell polarity. Nature 447: 330-333, 2007.

NCBI AND UNIPROT INFORMATION

NCBI GenBank Nucleotide #

[NM_001039469](#)

Molecular Weight

~114kDa

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/mark2-recombinant-protein-sf9-wb-western-blot-kinase-assay/73679>