

PRKACA (L206R) Recombinant Protein | PRKACA recombinant protein

PRKACA (L206R), Active

CATALOG # AAA73848	HOST Sf9 insect cells	APPLICATIONS Kinase Assay
------------------------------	---------------------------------	-------------------------------------


PRODUCT SYNONYMS

PRKACA (L206R); Active; PKACA; PPNAD4; PRKACA recombinant protein

PRODUCT SPECIFICATIONS


Host	Sf9 insect cells
Form/Format	Recombinant protein stored in 50mM Tris-HCl, pH7.5, 300mM NaCl, 10mM glutathione, 0.1mM EDTA, 0.25mM DTT, 0.1mM PMSF, 25% glycerol.
Applicable Applications for PRKACA recombinant protein	Kinase Assay
Tag	GST
Species	Human
Dry Ice Note	The product may be shipped with dry ice, and an additional fee may be added to your shipping cost.
Preparation and Storage	Store product at -70 degree C. For optimal storage, aliquot target into smaller quantities after centrifugation and store at recommended temperature. For most favorable performance, avoid repeated handling and multiple freeze/thaw cycles.

IMAGES



SDS-PAGE

Purity: Sample Purity Data. For specific information on a given lot, see related technical data sheet.



Activity

Specific Activity: Sample Activity Plot. For specific information on a given lot, see related technical data sheet.

ADDITIONAL INFORMATION

Related Product Information for PRKACA recombinant protein	Recombinant human PRKACA (1-351) was expressed by baculovirus in Sf9 insect cells using an N-terminal GST tag. The gene accession number is NM_002730. The catalytic subunit C-alpha of PRKACA is a member of the Ser/Thr protein kinase family and has been assigned to chromosome region 19p13.1 (1). Null mutation in PRKACA leads to early postnatal lethality in the majority of C-alpha knockout mice. Surprisingly, a small percentage of C-alpha knockout mice, although runted, survived to adulthood. In these animals, compensatory increases in C-beta levels occurred in brain whereas many tissues, including skeletal muscle, heart, and sperm, contained less than 10% of the normal PKA activity (2).
Product Categories/Family for PRKACA recombinant protein	Mutant Kinases; Cancer; Cardiovascular Disease

References

1. Tasken, K. et al: The gene encoding the catalytic subunit C-alpha of cAMP-dependent protein kinase (locus PRKACA) localizes to human chromosome region 19p13.1. Genomics 36: 535-538, 1996.
2. Skälhegg, BS. Et al: Mutation of the C-alpha subunit of PKA leads to growth retardation and sperm dysfunction. Molec. Endocr. 16: 630-639, 2002.

NCBI AND UNIPROT INFORMATION

NCBI GenBank Nucleotide #

[NM_002730](#)

Molecular Weight

65kDa

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/prkaca-l206r-recombinant-protein-sf9-kinase-assay/73848>