

RIPK3 recombinant protein

RIPK3, Active

CATALOG # AAA73920	HOST Sf9 insect cells	APPLICATIONS Western Blot
------------------------------	---------------------------------	-------------------------------------


PRODUCT SYNONYMS

RIPK3; Active; RIP3; RIPK3 recombinant protein

PRODUCT SPECIFICATIONS


Host	Sf9 insect cells
Form/Format	Recombinant protein stored in 50mM Tris-HCl, pH7.5, 150mM NaCl, 10mM glutathione, 0.1mM EDTA, 0.25mM DTT, 0.1mM PMSF, 25% glycerol.
Applicable Applications for RIPK3 recombinant protein	WB (Western Blot), Kinase Assay
Tag	GST
Species	Human
Dry Ice Note	The product may be shipped with dry ice, and an additional fee may be added to your shipping cost.
Preparation and Storage	Store product at -70 degree C. For optimal storage, aliquot target into smaller quantities after centrifugation and store at recommended temperature. For most favorable performance, avoid repeated handling and multiple freeze/thaw cycles.

IMAGES



SDS-PAGE

Purity: Sample Purity Data. For specific information on a given lot, see related technical data sheet.



Activity

Specific Activity: Sample Kinase Activity Plot. For specific information on a given lot, see related technical data sheet.

ADDITIONAL INFORMATION

Related Product Information for RIPK3 recombinant protein	Recombinant full-length human RIPK3 was expressed by baculovirus in Sf9 insect cells using an N-terminal GST tag. The RIPK3 gene accession number is NM_006871. RIPK3 is a member of the receptor-interacting protein (RIP) family of serine/threonine protein kinases, which contains a C-terminal domain unique from other RIP family members. RIPK3 is predominantly localized to the cytoplasm, and specifically, are colocalized in the mitochondrion which can undergo nucleocytoplasmic shuttling dependent on novel nuclear localization and export signals (1). RIPK3 is a component of the tumor necrosis factor (TNF) receptor-I signaling complex, and can induce apoptosis and weakly activate the NF-kappaB transcription factor and caspase-8 prevents RIPK3-dependent necrosis without inducing apoptosis by functioning in a proteolytically active complex with CFLAR and that this complex is required for the protective function (2).
Product Categories/Family for RIPK3 recombinant protein	Protein Kinases; Apoptosis/Autophagy; Cancer; Inflammation; JAK/STAT Pathway; JNK/SAPK Pathway; p38 Pathway; Ser/Thr Kinases

References

1. Kasof, G. M.et.al: The RIP-like kinase, RIP3, induces apoptosis and NF-kappa-B nuclear translocation and localizes to mitochondria. FEBS Lett. 473: 285-291, 2000.
2. Oberst, A. et.al: Catalytic activity of the caspase-8-FLIP(L) complex inhibits RIPK3-dependent necrosis. Nature 471: 363-367, 2011.

NCBI AND UNIPROT INFORMATION

NCBI GenBank Nucleotide #

[NM_006871](#)

Molecular Weight

~96kDa

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/ripk3-recombinant-protein-sf9-wb-western-blot-kinase-assay/73920>