

Goat anti Human IgE (epsilon chain) (HRP) Secondary Antibody | IgE secondary antibody

Goat anti Human IgE (epsilon chain) (HRP)

CATALOG # AAA75527	HOST Goat	APPLICATIONS Immunohistochemistry, Western Blot, ELISA
------------------------------	---------------------	--

PRODUCT SYNONYMS

Goat anti Human IgE (epsilon chain) (HRP); Goat anti Human Immunoglobulin E (epsilon chain) secondary (HRP); IgE secondary antibody

PRODUCT SPECIFICATIONS

Host	Goat
Form/Format	1.0mg/ml lyophilized from 10mM Sodium Phosphate, pH 7.2 with 150mM Sodium Chloride, with 1% BSA and 0.1% ProClin 150 (Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase).
Concentration	1.0 mg/ml, E ¹ % at 280 nm = 13.0 (prior to lyophilization) (varies by lot)
Applicable Applications for IgE secondary antibody	IHC (Immunohistochemistry), WB (Western Blot), ELISA
Cross-Reactivity	Reacts with heavy (Sigma) chains on Human IgE. Does not react with light chains on all Human immunoglobulins, or non-immunoglobulin Human serum proteins.
Immunogen	Purified Human IgE, ϵ chain
Reconstitution	Rehydrate with 1.1 ml of deionized water. Allow reconstituted product at room temperature at least 30 minutes prior to dilution.
Preparation and Storage	Upon receipt store lyophilized product at 2-8°C. After reconstitution, dilute with an equal volume of glycerol and store at -20°C. Prepare working dilution prior to use and then discard

ADDITIONAL INFORMATION

Product Categories/Family for IgE secondary antibody	Immunology; Conjugated Secondary Antibodies
--	---

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/goat-secondary-antibody/goat-anti-human-ige-epsilon-chain-hrp/75527>