

Goat anti Swine IgG (Fc specific) Secondary Antibody

Goat anti Swine IgG (Fc specific)

CATALOG # AAA77575	HOST Goat	REACTIVITY Swine
------------------------------	---------------------	----------------------------

PRODUCT SYNONYMS

Goat anti Swine IgG (Fc specific); Polyclonal Secondary Antibody; Goat anti Swine IgG (Fc specific) secondary antibody

PRODUCT SPECIFICATIONS

Host	Goat
Reactivity	Swine
Clonality	Polyclonal
Specificity	Precipitating polyclonal Goat antiSerum to Swine IgG, Fc specific.
Source	Highly purified normal IgG isolated from pooled swine serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Product	Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added. Reconstitute the lyophilized antiserum by adding 1 ml sterile distilled water.
Cross reactivity	The antiSerum does not cross-react with any other component of the Swine immunoglobulin system. Inter-species cross-reactivity is a normal feature of antibodies to immunoglobulins, since Ig of different species frequently share antigenic determinants. Cross-reactivity of this antiSerum has not been tested in detail.
Preparation and Storage	The lyophilized antiserum is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. Dilutions may be prepared by adding phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the antiserum. Diluted antiserum should be stored at +4°C, not refrozen, and preferably used the same day.

ADDITIONAL INFORMATION

Related Product Information for Goat anti Swine IgG (Fc specific) secondary antibody	The reactivity of the antiserum is restricted to the Fc part of the IgG molecule. In immunoelectrophoresis and radial immunodiffusion, using various antiserum concentrations against normal swine serum a single precipitin line is obtained which shows a reaction of identity with the precipitin line obtained with purified IgG. No precipitation reaction is obtained with purified IgA, IgM, and IgG/Fab fragments. In precipitating techniques as immunoelectrophoresis and radial immunodiffusion to identify the presence of IgG in swine serum or other body fluids or to determine its concentration. To prepare an immunoadsorbent for the purification of swine IgG from serum or plasma. This antiserum is not intended for use in non-precipitating antibody- binding or other highly sensitive assays. For such use labelled reagents, meeting the necessary specificity and performance requirements are available.
Product Categories/Family for Goat anti Swine IgG (Fc specific) secondary antibody	Antisera

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.