

Goat Anti-Mouse IgM+IgG+IgA(H+L) Secondary Antibody

Goat Anti-Mouse Ig, Human ads-FITC

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA78761	Goat	Mouse	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Immunoblot, Western Blot, ELISA

PRODUCT SYNONYMS

Goat Anti-Mouse IgM+IgG+IgA(H+L); Polyclonal Secondary Antibody; Goat Anti-Mouse Ig; Human ads-FITC; Human ads - FITC; Goat Anti-Mouse IgM+IgG+IgA(H+L) secondary antibody

PRODUCT SPECIFICATIONS

Host	Goat
Reactivity	Mouse
Clonality	Polyclonal
Isotype	IgG
Specificity	Reacts with the heavy and light chains of mouse IgG ₁ , IgG _{2a} , IgG _{2b} , IgG _{2c} , IgG ₃ , IgM, and IgA
Purity/Purification	Affinity chromatography on pooled mouse IgG,IgM,IgA covalently linked to agarose
Form/Format	The fluorescein (FITC) conjugates, are supplied as 1.0 mg in 1.0 mL PBS/NaN ₃ .
Applicable Applications for Goat Anti-Mouse IgM+IgG+IgA(H+L) secondary antibody	IS (Immunostaining), IB (Immunoblot), ELISA
Source	Pooled antisera from goats hyperimmunized with mouse IgG, IgM and IgA
Cross Absorption	Human immunoglobulins and pooled sera; may react with immunoglobulins from other species.
Preparation and Storage	Store at 2°C to 8°C. Protect fluorochrome-conjugated forms from light. Reagents are stable for the period shown on the label if stored as directed.

ADDITIONAL INFORMATION

Product Categories/Family for Goat Anti-Mouse IgM+IgG+IgA(H+L) secondary antibody	Goat Anti-Mouse IgM+IgG+IgA (H+L)
---	-----------------------------------

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/goat-anti-mouse-igmiggigahl-secondary-antibody-immunostaining-immunoblot-elisa/78761>