

Goat Anti-Porcine IgG (H+L) Secondary Antibody

Goat Anti-Porcine IgG (H+L)-HRP

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA78830	Goat	Porcine(may react with immunoglobulins from other species and the light chains of other porcine immunoglobulins)	Western Blot, Immunocytochemistry, Immunohistochemistry, Flow Cytometry, Functional Assay, ELISA

PRODUCT SYNONYMS

Goat Anti-Porcine IgG (H+L); Polyclonal Secondary Antibody; Goat Anti-Porcine IgG (H+L)-HRP; Goat Anti-Porcine IgG(H+L) - HRP; Goat Anti-Porcine IgG (H+L) secondary antibody

PRODUCT SPECIFICATIONS

Host	Goat
Reactivity	Porcine (may react with immunoglobulins from other species and the light chains of other porcine immunoglobulins)
Clonality	Polyclonal
Isotype	IgG
Specificity	Reacts with the heavy and light chains of porcine IgG
Purity/Purification	Affinity chromatography on porcine IgG covalently linked to agarose.
Form/Format	HRP conjugated supplied as 1.0 mL in a stock solution of 50% glycerol/50% PBS, pH7.4. No preservative added.
Applicable Applications for Goat Anti-Porcine IgG (H+L) secondary antibody	WB (Western Blot), ICC (Immunocytochemistry), IHC (Immunohistochemistry), FCM/FACS (Flow Cytometry), FLISA, ELISA
Source	Pooled antisera from goats hyperimmunized with porcine IgG.
Preparation and Storage	Store at 2 to 8°C or long-term at -20°C

ADDITIONAL INFORMATION

Product Categories/Family for Goat Anti-Porcine IgG (H+L) secondary antibody	Goat Anti-Porcine IgG (H+L)
--	-----------------------------

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/goat-anti-porcine-igg-hl-secondary-antibody-western-blot-immunocytochemistry-immunohistochemistry-flow-cytometry-elisa/78830>