

Goat Anti-Feline IgG (H+L) Secondary Antibody

Goat Anti-Feline IgG (H+L)-HRP

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA78835	Goat	Feline	Western Blot, Immunohistochemistry, Immunofluorescence, Immunoblot, Western Blot, ELISA


PRODUCT SYNONYMS

Goat Anti-Feline IgG (H+L); Polyclonal Secondary Antibody; Goat Anti-Feline IgG (H+L)-HRP; Goat Anti-Feline IgG(H+L) - HRP; Goat Anti-Feline IgG (H+L) secondary antibody

PRODUCT SPECIFICATIONS

Host	Goat
Reactivity	Feline
Clonality	Polyclonal
Isotype	IgG
Specificity	Reacts with the heavy and light chains of feline IgG as demonstrated by ELISA or FLISA. May also react with the light chains of other feline immunoglobulins.
Purity/Purification	Affinity chromatography on feline IgG covalently linked to agarose.
Form/Format	Horseradish Peroxidase (HRP) Conjugate is supplied as 1.0 mL in a stock solution of 50% glycerol, pH7.4.
Applicable Applications for Goat Anti-Feline IgG (H+L) secondary antibody	WB (Western Blot), IHC (Immunohistochemistry), IF (Immunofluorescence), IB (Immunoblot), FLISA, ELISA
Source	Pooled antisera from goats hyperimmunized with feline IgG.
Cross Absorption	None; may react with immunoglobulins for each other species and the light chains of other feline immunoglobulins.
Warning	Reagents contain sodium azide. Sodium Azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.
Preparation and Storage	Store at 2-8 degree C or long-term at -20 degree C.

IMAGES



Application Data

FLISA plate was coated with purified feline IgG. Immunoglobulin was detected with Goat Anti-Feline IgG(H+L)-FITC

ADDITIONAL INFORMATION

Product Categories/Family for Goat Anti-Feline IgG (H+L) secondary antibody	Goat Anti-Feline IgG (H+L)
---	----------------------------

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/goat-anti-feline-igg-hl-secondary-antibody-western-blot-immunohistochemistry-immunofluorescence-immunoblot-elisa/78835>