

VEGF-C active protein

Human VEGF-C

CATALOG # AAA79143	HOST Insect Cells	REACTIVITY Human
------------------------------	-----------------------------	----------------------------


PRODUCT SYNONYMS

VEGF-C; Human VEGF-C; Recombinant Human Vascular Endothelial Growth Factor-C; VRP; Flt4-L; VEGF-related protein; VEGF-C active protein


PRODUCT SPECIFICATIONS

Host	Insect Cells
Reactivity	Human
Purity/Purification	> 90% by SDS-PAGE & silver stain
Form/Format	Lyophilized
Sequence	N-Terminal Sequence:DPTEETI
Sequence Length	121
N Terminal Sequence	DPTEETI
Buffer	Water
Preparation and Storage	Lyophilized samples are stable for more than six months at -20 degree C to -70 degree C. Reconstituted VEGF-C should be stored in working aliquots at -20 degree C. Avoid repeated freeze-thaw cycles.

IMAGES



Application Data



Application Data

ADDITIONAL INFORMATION

Related Product Information for VEGF-C active protein	VEGF-C, also known as Vascular Endothelial Growth Factor Related Protein (VRP), is a recently discovered VEGF growth factor family member that is most closely related to VEGF-D. The human VEGF-C cDNA encodes a pre-pro-protein of 416 amino acids residues. It is almost identical to the mouse VEGF-C protein. Similar to VEGF-D, VEGF-C has a VEGF homology domain spanning the middle third of the precursor molecule and long N- and C-terminal extensions. In adults, VEGF-C is highly expressed in heart, placenta, ovary and small intestine. Recombinant human VEGF-C, lacking the N- and C-terminal extensions and containing only the middle VEGF homology domain, forms primarily non-covalently linked dimers. This protein is a ligand for both VEGFR-2/KDR and VEGFR-3/FLT-4. Since VEGFR-3 is strongly expressed in lymphatic endothelial cells, it has been postulated that VEGF-C is involved in the regulation of the growth and/or differentiation of lymphatic endothelium. Although recombinant human VEGF-C is also a mitogen for vascular endothelial cells, it is much less potent than VEGF-A. The recombinant human VEGF-C contains 121 amino acids residues and was fused to a His-tag (6x His) at the C-terminal end. As a result of glycosylation VEGF-C migrates as an 18-24 kDa protein in SDS-PAGE under reducing conditions.
Product Categories/Family for VEGF-C active protein	Cytokines & Growth Factors
References	1. Joukov et al., EMBO J 15:290, 1996 2. Olofsson et al., Curr Opin Biotech 10:528, 1999 3. Kukk et al., Development 122:3829, 1996

NCBI AND UNIPROT INFORMATION

NCBI GI #	4885653
NCBI GeneID	7424
NCBI Accession #	NP_005420.1
NCBI GenBank Nucleotide #	NM_005429.2
UniProt Accession #	P49767; B2R9Q8
Molecular Weight	18-24 kDa
NCBI Official Full Name	vascular endothelial growth factor C preproprotein
NCBI Official Synonym Full Names	vascular endothelial growth factor C
NCBI Official Symbol	VEGFC
NCBI Official Synonym Symbols	VRP; Flt4-L
NCBI Protein Information	vascular endothelial growth factor C; VEGF-C; FLT4 ligand DHM; vascular endothelial growth factor-related protein
UniProt Protein Name	Vascular endothelial growth factor C
UniProt Gene Name	VEGFC
UniProt Synonym Gene Names	VEGF-C; Flt4-L; VRP
UniProt Entry Name	VEGFC_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.