

Human noggin ELISA Kit | NOG elisa kit

Human Noggin, NOG ELISA Kit

CATALOG #
AAA80883

REACTIVITY
Human


PRODUCT SYNONYMS

noggin; Human Noggin; NOG ELISA Kit; Human Noggin (NOG) ELISA kit; SYM1; SYNS1; symphalangism 1 (proximal) ; NOG elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of human NOG. No significant cross-reactivity or interference between human NOG and analogues was observed.
Sequence Length	232
Samples	Serum, plasma
Assay Type	Sandwich
Detection Range	1.56 ng/ml-100 ng/ml.
Sensitivity	The minimum detectable dose of human NOG is typically less than 0.39 ng/ml. The sensitivity of this assay, or Lower Limit of Detection (LLD) was defined as the lowest protein concentration that could be differentiated from zero. It was determined the mean O.D value of 20 replicates of the zero standard added by their three standard deviations.
Intra-assay Precision	Intra-assay Precision (Precision within an assay): CV%<8%. Three samples of known concentration were tested twenty times on one plate to assess.
Inter-assay Precision	Inter-assay Precision (Precision between assays): CV%<10%. Three samples of known concentration were tested in twenty assays to assess.
Preparation and Storage	Unopened test kits should be stored at 2 to 8 degree C upon receipt. Please refer to pdf manual for further storage instructions.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for NOG elisa kit	Principle of the assay: This assay employs the quantitative sandwich enzyme immunoassay technique. Antibody specific for NOG has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and any NOG present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for NOG is added to the wells. After washing, avidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of NOG bound in the initial step. The color development is stopped and the intensity of the color is measured.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	4885523
NCBI GeneID	9241
NCBI Accession #	NP_005441.1
NCBI GenBank Nucleotide #	NM_005450.4
UniProt Accession #	Q13253
Molecular Weight	25,774 Da
NCBI Official Full Name	noggin
NCBI Official Synonym Full Names	noggin
NCBI Official Symbol	NOG
NCBI Official Synonym Symbols	SYM1; SYNS1
NCBI Protein Information	noggin; symphalangism 1 (proximal)
UniProt Protein Name	Noggin
UniProt Gene Name	NOG
UniProt Entry Name	NOGG_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.