

Mouse Interleukin 12 ELISA Kit | IL-12/P70 elisa kit

Mouse Interleukin 12 ELISA Kit

CATALOG #
AAA85282

REACTIVITY
Mouse


PRODUCT SYNONYMS

Interleukin 12; Mouse Interleukin 12 ELISA Kit; Interleukin 12 (Mouse); IL-12/P70 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Assay Type	Sandwich
Preparation and Storage	Store all reagents at 2-8 degree C

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for IL-12/P70 elisa kit

Intended Uses: This IL-12 ELISA kit is intended Laboratory for research use only. The Stop Solution changes the color from blue to yellow and the intensity of the color is measured at 450 nm using a spectrophotometer. In order to measure the concentration of IL-12 in the sample, this IL-12 ELISA Kit includes a set of calibration standards. The calibration standards are assayed at the same time as the samples and allow the operator to produce a standard curve of Optical Density versus IL-12 concentration. The concentration of IL-12 in the samples is then determined by comparing the O.D. of the samples to the standard curve.

Principle of the Assay: This IL-12 enzyme linked immunosorbent assay applies a technique called a quantitative sandwich immunoassay. The microtiter plate provided in this kit has been pre-coated with a monoclonal antibody specific for IL-12. Standards or samples are then added to the microtiter plate wells and IL-12 if present, will bind to the antibody pre-coated wells. In order to quantitatively determine the amount of IL-12 present in the sample, a standardized preparation of horseradish peroxidase (HRP)-conjugated polyclonal antibody, specific for IL-12 are added to each well to "sandwich" the IL-12 immobilized on the plate. The microtiter plate undergoes incubation, and then the wells are thoroughly washed to remove all unbound components. Next, A and B substrate solution is added to each well. The enzyme (HRP) and substrate are allowed to react over a short incubation period. Only those wells that contain IL-12 and enzyme-conjugated antibody will exhibit a change in colour. The enzyme-substrate reaction is terminated by the addition of a sulphuric acid solution and the colour change is measured spectrophotometrically at a wavelength of 450 nm.

NCBI AND UNIPROT INFORMATION

NCBI GI #	4323579
NCBI GeneID	3592
UniProt Accession #	P29459; Q96QZ1
Molecular Weight	24,874 Da
NCBI Official Full Name	interleukin 12
NCBI Official Synonym Full Names	interleukin 12A (natural killer cell stimulatory factor 1, cytotoxic lymphocyte maturation factor 1, p35)
NCBI Official Symbol	IL12A
NCBI Official Synonym Symbols	P35; CLMF; NFSK; NKSF1; IL-12A
NCBI Protein Information	interleukin-12 subunit alpha; CLMF p35; IL35 subunit; IL-12 subunit p35; IL-12, subunit p35; interleukin 12, p35; interleukin-12 alpha chain; NF cell stimulatory factor chain 1; NK cell stimulatory factor chain 1; cytotoxic lymphocyte maturation factor 1, p35; cytotoxic lymphocyte maturation factor 35 kDa subunit; natural killer cell stimulatory factor 1, 35 kD subunit
UniProt Protein Name	Interleukin-12 subunit alpha
UniProt Gene Name	IL12A
UniProt Synonym Gene Names	NKSF1; IL-12A; CLMF p35; NKSF1
UniProt Entry Name	IL12A_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.