

Human Vanillylmandelic acid ELISA Kit | VMA elisa kit

Human Vanillylmandelic acid ELISA Kit

CATALOG #
AAA86405

REACTIVITY
Human


PRODUCT SYNONYMS

Vanillylmandelic acid; Human Vanillylmandelic acid ELISA Kit; VMA elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of VMA. No significant cross-reactivity or interference between VMA and analogues was observed. NOTE: Limited by current skills and knowledge, it is impossible for us to complete the cross-reactivity detection between VMA and all the analogues, therefore, cross reaction may still exist in some cases.
Samples	Serum, plasma, cell culture supernatants, body fluid and tissue homogenate
Assay Type	Quantitative Competitive
Sensitivity	0.1 mg/L
Preparation and Storage	Store all reagents at 2-8 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for VMA elisa kit	Intended Uses: This VMA ELISA kit is a 1.5 hour solid-phase ELISA designed for the quantitative determination of Human VMA. This ELISA kit for research use only, not for therapeutic or test applications! Principle of the Assay: VMA ELISA kit applies the competitive enzyme immunoassay technique utilizing an anti-VMA antibody and an VMA-HRP conjugate. The assay sample and buffer are incubated together with VMA-HRP conjugate in pre-coated plate for one hour. After the incubation period, the wells are decanted and washed five times. The wells are then incubated with a substrate for HRP enzyme. The product of the enzyme-substrate reaction forms a blue colored complex. Finally, a stop solution is added to stop the reaction, which will then turn the solution yellow. The intensity of color is measured spectrophotometrically at 450nm in a microplate reader. The intensity of the color is inversely proportional to the VMA concentration since VMA from samples and VMA-HRP conjugate compete for the anti-VMA antibody binding site. Since the number of sites is limited, as more sites are occupied by VMA from the sample, fewer sites are left to bind VMA-HRP conjugate. A standard curve is plotted relating the intensity of the color (O.D.) to the concentration of standards. The VMA concentration in each sample is interpolated from this standard curve.
---	---

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/vma-human-elisa-kits/vanillylmandelic-acid/86405>